



**NNS SHOW GOAT M20 TXT  
MEDICATED**

For the prevention of coccidiosis caused by Eimeria crandalis, Eimeria christenseni and Eimeria ninakohlyakimovae.

**ACTIVE DRUG INGREDIENT**

Monensin (as monensin sodium) ..... 20.0 g/ton

**GUARANTEED ANALYSIS**

Crude Protein not less than 15.90 %

\* This includes not more than 1.5 % equivalent crude protein from non-protein nitrogen

Crude Fat			not less than	3.06 %
Crude Fiber			not more than	3.33 %
Calcium	not less than	1.57	not more than	1.99 %
Phosphorus			not less than	0.35 %
Salt	not less than	0.71	not more than	0.90 %
Copper	not less than	24.4	not more than	24.9 ppm
Selenium			not less than	0.4 ppm
Vitamin A			not less than	5,030 IU/lb

**INGREDIENTS**

Grain Products, Processed Grain By-Products, Plant Protein Products, Calcium Carbonate, Ammonium Chloride, Salt, Molasses Products, Vitamin E Supplement, Urea Formaldehyde Condensation Polymer, Vitamin A Supplement, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Vitamin D3 Supplement, Ethylenediamine Dihydrochloride.

**FEEDING DIRECTIONS**

For growing and finishing goats. Gradually acclimate goats to feed over a 4-6-week period to avoid digestive upsets. After acclimation period, feed up to 3% of goat body weight daily split into multiple feedings. Always feed in combination with forage. Always provide fresh, clean water

**CAUTION:** Do not allow horses or other equines access to feeds containing Monensin. Monensin medicated goat and cattle feeds are safe for use in goats and cattle only. Consumption by unapproved species may result in toxic reactions.

**LIMITATIONS:** Feed continuously. Do not feed to lactating goats.

UniBond(Urea Formaldehyde Condensation Polymer)Pellet Binder "Approved For Use In All Animal Feeds"

FS4197

**Manufactured By:  
Agfinity, Inc.**

**260 Factory Road - Eaton, CO 80615  
Net Weight Bulk-or-50 lb (22.67 Kg)**

**MEDICATED**